

DAFTAR PUSTAKA

- Agati, G., Azzarello, E., Pollastri, S., and Tattini, M. (2012). Flavonoids as Antioxidants in Plants: Location and Functional Significance. *Plant Science*, 196: 67-76.
- Aida, M., Shinomiya, K., Matsuzawa, K., Hano, Y., Nomura, T. (1994). Constituents of The Moraceae Plants. 19. Artonins Q, R, S, T, and U, Five New Isoprenylated Phenols from The Bark of *Artocarpus heterophyllus* Lamk. *Heterocycle*, 39: 847-858.
- Aisyah, S. (2008). *Bahan Ajar Penentuan Struktur Senyawa Organik*. Bandung: FPMIPA UPI.
- Arung, E.T., Yoshikawa, K., Shimizu, K., Kondo, R. (2010). Isoprenoid-Substituted Flavonoids from Wood of *Artocarpus heterophyllus* on B16 Melanoma Cells: Cytotoxicity and Structural Criteria. *Fitoterapia*, 81(2):120-123.
- Attioua, B., Lagnika, L., Yeo, D., Antheaume, C., Kaiser, M., Weniger, B., Lobstein, A., Sénécheau, C.V. (2011). *In Vitro* Antiplasmodial and Antileishmanial Activities of Flavonoids from *Anogeissus leiocarpus* (Combretaceae). *International Journal of Pharmaceutical Sciences Review and Research*, 11(2): 1-6.
- Barliana, M.I., Diantini, A., Abdulah, R. (2012). *Aktivitas Tanaman Asli Indonesia Puspa (Schima wallichii) sebagai Senyawa Antimalaria Baru*. Bandung: Universitas Padjadjaran.
- Barzinji, A.K.R., Nasher, A.K., Mothana, R.A. dan Al-Hamadi, M.S. (2014). *In Vitro* Antimalarial Activity of Selected Yemeni Plants Used in Traditional Medicine. *International Journal of Medicinal Plants*, 107:526-535.
- Bhat, G.P. dan Surolia, N. (2001). *In Vitro* Antimalarial Activity of Extracts of Three Plants Used in The Traditional Medicine of India. *The American Society of Tropical Medicine and Hygiene*, 65(4): 304-308.
- Boonphong, S., Baramee, A., Kittakoop, P. Dan Puangsombat, P. (2007). Antitubercular and Antiplasmodial Prenylated Flavones from The Roots of *Artocarpus altilis*. *Chiang Mai J. Sci.*, 34(3): 339-344.
- Brunetti, C., Ferdinando, M.D., Fini, A., Pollastri, S., and Tantini, M. (2013). Flavonoids as Antioxidants and Developmental Regulators: Relative

- Significance in Plants and Humans. *International Journal of Molecular Science*, 14: 3540-3555.
- Candrika. (2006). Hypoglycaemic Action Of The Flavanoid Fraction of *Artocarpus heterophyllus* Leaf. *Afr. J. Trad. CAM*, 3(2): 42-50.

- Cannel, R.J.P. (1998). *How to Approach the Isolation of A Natural Product dalam Natural Products Isolation*. New Jersey: Humana Press.
- Cartea ME, Francisco M, Lema M, Soengas P, Velasco P. (2010). Resistance of cabbage (*Brassica oleracea capitata* Group) Crops to *Mamestra brassicae*. *Journal of Economic Entomology*, 103(5): 1866- 1874.
- Chintamunnee, V. Dan Mahomoodally, M.F. (2012). Herbal Medicine Commonly Used Against Non-Communicable Diseases in The Tropical Island of Mauritius. *Jornal of Herbal Medicine*, 2(4): 113-125.
- Day, R. A. Jr. dan Underwood, A. L. (2002). *Analisis Kimia Kualitatif Edisi Keenam*. Jakarta: Erlangga.
- Devasagayam, T.P.A., Tilak, J.C., Singhal, R. (2001). *Functional Foods in India; History And Scope in Angiogenesis*. In: Losso, J.N., Shahidi, F., Bagchi, D., Editor: *Functional and Medicinal Foods*. New York: New York In press.
- Di, X., Wang, S., Wang, B., Liu, Y., Yuan, H., Lou, H., Wang, X. (2013). New Phenolic Compounds from the Twigs of *Artocarpus heterophyllus*. *Drug Discoveries and Therapeutics*, 7(1): 24-28.
- Elfahmi, Woeedenbag, H.J., Kayser, O. (2014). Jamu: Indonesia Traditional Herbal Medicine Towards Rational Phytopharmacological Use. *Journal of Herbal Medicine*, 4(2): 51-73
- Ersam, T. (2001). *Senyawa Kimia Makro Molekul Beberapa Tumbuhan Artocarpus Hutan Tropika Sumatera Barat*. Bandung: Disertasi ITB
- Ersam, T. (2001). *Senyawa Kimia Makromolekul Beberapa Tumbuhan Artocarpus Hutan Tropika Sumatera Barat*. Bandung: ITB.
- Fang, S.C., Hsu, C.L., Dan Yen, G.C. (2008). Anti-Inflammatory Effects Of Phenolic Compounds Isolated From The Fruits Of *Artocarpus Heterophyllus*. *Journal Agriculture Food Chem*, 56 (12), 4463-4468.
- Fidock DA, Rosenthal PJ, Croft SL, Brun R, Nwaka S. (2004). Antimalarial Drug Discovery: Efficacy Models for Compound Screening Review. *Nature*, 3: 509-520.

- Ghisalberti, E.L. (2008). *Detection and Isolation of Bioactive Natural Products dalam Bioactive Natural Product: Detection, Isolation and Structural Determination 2nd Edition*. New York: CRC Press.
- Gupta, D., Mann, S., Sood, A., Gupta, R.K. (2011). Phytochemical, Nutritional And Antioxidant Activity Evaluation Of Seeds Of Jackfruit (*Artocarpus heterophyllus* Lamk.). *International Journal Of Pharma And Bio Science*, 2(4),336-345.
- Han, Jang-na. (2012). Treatment of Influenza: Chinese Medicine Vs. Western Medicine. *Journal of Thoracic Disease*, 4(1):10-11.
- Hano, Y., Aida, M., Nomura, T. (1990). Two New Natural Diels-Alder Type Adducts from the Root Bark of *Artocarpus heterophyllus*. *Journal of Natural Product*, 53: 391-395.
- Hawaii Tropical Botanical Garden. (2014). *Hawaii Tropical Botanical Garden- Artocarpus heterophyllus*. [Online]. Tersedia: <http://www.htbg.com/Moraceae/ART-001-1-20-004/>. [18 September 2014].
- Herlina, T., Supratman, U., Urbanas, A., Sutardjo, S., Abdullah, N.R., Hayashi, H. (2011). Triterpenoid Pentacyclic Antimalarial Activity from the Leaves of *Erythrina variegata*. *Jurnal ILMU DASAR*, 12:161-166.
- Heyne, K. (1987). Tumbuhan Berguna Indonesia Jilid II. Jakarta: Badan Litbang Kehutanan.
- Hostettmann,K., Hostettmann, M., and Marston, A. (1995). *Cara Kromatografi Preparatif*. Bandung: ITB Press.
- Ichino C, et al. (2006). Antimalarial Activity Of Biflavonoids from *Ochna integerrima*. *Planta Med*, 72: 611–614.
- Ito J, Ghosh A, Moreira LA, Wimmer EA, Jacobs-Lorena M. (2002). Transgenic Anopheline Mosquitoes Impaired In Transmission Of A Malaria Parasite, *Nature*. 417: 387-398.
- Kaur, S dan Mondal P. (2014). **Study of Total Phenolic and Flavonoid Content, Antioxidant Activity and Antimicrobial Properties of Medicinal Plants.** *Journal of Microbiology and Experimentation*, 1(1): 1-6.

- Kementrian Kesehatan RI. (2011). *Buletin-Jendela Data dan Informasi Kesehatan: Epidemiologi Malaria di Indonesia*. Jakarta: Kementrian Kesehatan Republik Indonesia.
- Khaomek P et al. (2008). *In Vitro Antimalarial Activity Of Prenylated Flavonoids from Erythrina fusca*, J Nat Med, 62: 217–220.
- Kijjoa, A., Cidade, H.M., Pinto, M.M.M., Gonzalez, M.J.T.G., Anantachoke, C., Gedris, T.E., Herz, W. (1996). Prenylflavonoids from *Artocarpus elasticus*. *Phytochemistry*, 43: 691-694.
- Kim, Y.C., Kim, H.S., Wataya, Y., Sohn, D.H., Kang, T.H., Kim, M.S., Kim, Y.M., Lee, G.M., Chang, J.D. dan Park, H. (2004). Antimalarial Activity of Lavandulyl Flavanones Isolated from The Roots of *Sophora flavescens*. *Biological and Pharmaceutical Bulletin*, 27: 748-750
- Kristanti, A.N., Aminah, N.S., Tanjung, M., Kurniadi, B. (2008). *Buku Ajar Fitokimia*. Surabaya: Airlangga University Press.
- Lemmens, Soerianegara. (1995). *Prosea (Plant Resource of South East Asia) Timber Trees : Minor Commercial Timber*. Bogor: Prosea Foundation.
- Lin, C., Lu, C., Huang, P. (1995). Flavonoids from *Artocarpus heterophyllus*. *Phytochemistry*, 39(6):1447-1451.
- Madhavi, D.L., Singhal, R.S., Kulkarni, P.R. (1985). *Technological Aspects of Food Antioxidants dalam D.L. Madhavi, S.S. Deshpande dan D.K. Salunkhe: Food Antioxidant, Technological, Toxicological and Health Perspectives*. Hongkong: Marcel Dekker Inc.
- Maslarova, N.V. Yanishlieva. (2001). *Inhibiting oxidation dalam Jan Pokorný, Nedyalka Yanislieva dan Michael Gordon: Antioxidants in food, Practical applications*. Cambridge: Woodhead Publishing Limited.
- Muhiedin, Fuad. (2008). *Efisiensi Proses Ekstraksi Oleoresin Lada Hitam dengan Metode Ekstraksi Multi Tahap*. Malang: Universitas Brawijaya.
- Mulja, M. dan Suharman. (1995). *Analisis Instrumental*. Surabaya: Airlangga University Press.

- Musthapa, I., Hakim, E.H., Juliawaty, L.D., Syah. Y.M., dan Achmad, S.A. (2010). Prenylated Flavones from Some Indonesian *Artocarpus* and Their Antimalarial Properties. *Medicinal Plants*, 2(2): 145-148.
- Nagavani, V., Rao, T.R. (2010). Evaluation of Antioxidant Potential and Qualitative Analysis of Major Polyphenols by RP-HPLC in *Nymphaea nouchali Burm* Flowers. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2(4): 98-104.
- Namdaung U *et al.* (2006). Bioactive Constituents Of The Root Bark Of *Artocarpus rigidus* Subsp. *Rigidus*. *Chem Pharm Bull*, 54: 1433–1436.
- Narasimhan, R., Toshihiko Osawa, Mitsuo Namiki and Shunro Kawakishi. (1988). Chemical Studies on Novel Rice Hull Antioxidants. 1. Isolation, Fractination, and Partial Characterization. *J. Agric. Food. Chem*, 37: 732-737.
- Nomura, T., Hano, Y., Aida, M. (1998). Isoprenoid Substituted Flavanoids from *Artocarpus* Plants (Moraceae). *Heterocycles*, 47: 1179-1205.
- Panthong, K., Tohdee, K., Towatana, N.H., Voravuthikunchai, S.P., Chusri, S. (2013). Prenylated Flavone from Roots of Hybrid Between *Artocarpus heterophyllus* and *Artocarpus integer* and Its Biological Activities. *Journal of The Brazilian Chemical Society*, 24(10): 1656-1661.
- Prohati. (2014). Detil Data *Artocarpus heterophyllus*. [Online]. Tersedia: <http://www.proseanet.org/prohati2/browser.php?docsid=243>. [18 September 2014].
- Radwan, M.M. *et al.* (2008). Non-Cannabinoid Constituents From A High Potency *Cannabis sativa* Variety. *Phytochemistry*, 69: 2627–2633.
- Rajalakshmi, D. dan S. Narasimhan. (1985). *Food Antioxidants: Sources and Methods of Evaluation dalam D.L. Madhavi: Food Antioxidant, Technological, Toxicological and Health Perspectives*. Hongkong: Marcel Dekker Inc.
- Sam, G.H., Mensah, M.L.K., Annan, K., Zahree, S. (2011). Plants Traditionally Used in Treating Malaria, Typhoid Fever and Piles in The Wa Municipality and Wa East (Funsi) District of The Upper West Region of Ghana. *Academic Journal*, 5(10): p3352.

- Santosa, C.M. dan Hertiani, T. (2005). Kandungan Senyawa Kimia dan Efek Ekstrak Air Daun Bangun-Bangun (*Coleus amboinicus* L) pada Aktivitas Fagositosis Netrofil Tikus Putih. *Maj Farm Indones*, 16:141-148.
- Sardjono, R.E. dan Dwiyanti, G. (2002). *Petunjuk Praktikum Kimia Organik II*. Bandung: FPMIPA UPI.
- Sarker, S.D., Latif, Z., and Gray, A.L. (2006). *Natural Product Isolation*. New Jersey: Humana Press.
- Sastrohamidjojo, H. (2002). *Kromatografi*. Yogyakarta: Liberty.
- Scholichah, V. (2012). Kualitas Mikrobiologi Jamu Gendong Jenis Kunir Asem yang Diproduksi di Kelurahan Merbung, Kecamatan Klaten Selatan, Kabupaten Klaten. *Jurnal Kesehatan Masyarakat*, 1(2): 504-513.
- Shanmugapriya, K., Saravana, P.S., Payal, H., Mohammed, S.P., Binnie, W. (2011). Antioxidant Activity, Total Phenolic and Flavonoid Contents of *Artocarpus heterophyllus* and *Manilkara zapota* Seeds and Its Reduction Potential. *International Journal of Pharmacy and Pharmaceutical Sciences*, 3:256-260.
- Skolnick, A.A. (1997). Old Chinese Herbal Medicine Used for Fever Yields Possible New Alzheimer Disease Theraphy. *The Journal of The American Medical Association*, 227(10):776.
- Soebagio, dkk. (2005). *Kimia Analitik II*. Malang: Universitas Negeri Malang (UM Press).
- Sudarmadji, S., Haryono B., dan Suhardi. (1989). *Analisa Bahan Makanan dan Pertanian*. Yogyakarta: Liberty.
- Susanto, Cornelius Eko. (2013). *Tinggi, Kasus Malaria di Indonesia*. [Online]. Tersedia:[http://www.metrotvnews.com/metronews/read/2013/04/23/3/148819/Tinggi-Kasus-Malaria-di-Indonesia-\[23-April-2013\]](http://www.metrotvnews.com/metronews/read/2013/04/23/3/148819/Tinggi-Kasus-Malaria-di-Indonesia-[23-April-2013]).
- Swantara, M.D., Darmayasa, I.B.G., dan Dewi, N.K.A.K. (2011). Uji Aktivitas Antibakteri Fraksi Kulit Batang Nangka. *Jurnal Kimia*, 5(1): 1-8.
- Tarigan, J.Br., Zuhra, C.F., and Sihotang,H. (2008). Skrining Fitokimia Tumbuhan yang Digunakan Oleh Pedagang Jamu Gendong untuk Merawat Kulit Wajah di Kecamatan Medan Baru. *Jurnal Biologi Sumatera*, 3(1):1-6.

- Trager, W., Jensen, J., 1976. Human Malaria Parasite in Continuous Culture. *Science*, 193: 673-675.
- Tran, L.Q., Tezuka, Y., Ueda, J., Nguyen, N.T., Maruyama, Y., Begum, K., Kim, H.S., Wataya, Y., Tran, Q.K., dan Kadota, S. (2003). *In Vitro* Antiplasmodial Activity of Antimalarial Medicinal Plants Used in Vietnamese Traditional Medicine. *Journal of Ethnopharmacology*, 86: 249-252.
- United States Department of Agriculture Natural Resources Conservation Service. (2014). *Artocarpus heterophyllus* Lam Jackfruit. [Online]. Tersedia: <http://plants.usda.gov/core/profile?symbol=arhe2>. [18 September 2014].
- Valdés, A.F.C., Martinez, J.M., Lizama, R.S., Gaitén, Y.G., Rodriguez, D.A. dan Payrol, J.A. (2010). *In Vitro* Antimalarial Activity and Cytotoxicity of Some Selected Cuban Medicinal Plants. *Rev.Inst. Med. Trop. Sao Paulo*, 52(4): 197-201.
- Venkataraman K. (1972). Wood phenolics in the chemotaxonomy of the Moraceae. *Phytochem*, 11: 1571-1586.
- Weniger B *et al.* (2004). A Bioactive Biflavonoid from *Campnosperma panamense*. *Fitoterapia*, 75: 764–767.
- White, P.J. and Y. Xing. (1954). *Antioxidants from Cereals and Legumes dalam Foreidoon Shahidi: Natural Antioxidants, Chemistry, Health Effect and Applications*. Illinois : AOCS Press.
- WHO Media Centre (2014). *Malaria*. [Online]. Tersedia: <http://www.who.int/mediacentre/factsheets/fs094/en/>. [21 Agustus 2014].
- Widyawaruyanti, A., Devi, A.P., Fatria, N., Tumewu,L., Tantular, I.S. dan Hafid, A.F. (2014). *In Vitro* Antimalarial Activity Screening of Several Indonesian Plants Using HRP2 Assay. *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(6): 125-128.
- Widyawaruyanti, A.S., Kalauni, S.K., Awale, S., Nindatu, M., Zaini, N.C., Syafruddin, D., Asih, P.B.S., Tezuka, Y. Dan Kadota, S. (2007). New Prenylated Flavones from *Artocarpus champeden*, and Their Antimalarial Activity *in vitro*. *Journal of Natural Medicines*, 61: 410-413.

- Yao Y, Vieira A (2011) Comparative Antioxidant Properties of *Citrus species*: Evidence for Potent, Non-Vitamin Antioxidants in *C. Aurantifolia*. *International Journal of Food, Nutrition and Public Health*, 4: 108-112.
- Yenjai, C., Prasanphen, K., Daodee, S., Wongpanich, V. Dan Kittakoop, P. (2004). Bioactive Flavonoids from *Kaempferia parviflora*. *Fitoterapia*, 75: 89-92.
- Zuhra, C.F., Tarigan, J.Br., and Sihotang, H. (2008). Aktivitas Antioksidan Senyawa Flavonoid dari Daun Katuk. *Jurnal Biologi Sumatera*, 3(1): 7-10.