

SPSS Data Pretest

Group Statistics

	Kecerdasan visual-spasial	N	Mean	Std. Deviation	Std. Error Mean
Pretest	kel eksperimen	15	26,13	1,685	,435
	kel kontrol	13	25,62	2,844	,789

Independent Samples Test

	Levene's Test for Equality of Variances	t-test for Equality of Means								
		F	Sig.	T	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Pretest	Equal variances assumed	2,030	,166	,596	26	,556	,518	,869	-1,269	2,305
	Equal variances not assumed			,575	18,908	,572	,518	,901	-1,368	2,404

Tests of Normality

		Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Pretest	kel eksperimen	,163	15	,200 [*]	,964	15	,755
	kel kontrol	,131	13	,200 [*]	,957	13	,712

Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
Pretest	Based on Mean	2,030	1	26	,166
	Based on Median	1,780	1	26	,194
	Based on Median and with adjusted df	1,780	1	19,038	,198
	Based on trimmed mean	1,949	1	26	,175

SPSS Data Posttest

Tests of Normality

	Kecerdasan visual-spasial	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Posttest	kel eksperimen	,273	15	,056	,646	15	,000
	kel kontrol	,203	13	,158	,943	13	,501

Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
Posttest	Based on Mean	,122	1	26	,729
	Based on Median	,036	1	26	,850
	Based on Median and with adjusted df	,036	1	21,192	,851
	Based on trimmed mean	,025	1	26	,875

Test of Homogeneity of Variance

	Levene Statistic	df1	df2	Sig.	
Posttest	Based on Mean	,122	1	26	,729
	Based on Median	,036	1	26	,850
	Based on Median and with adjusted df	,036	1	21,192	,851
	Based on trimmed mean	,025	1	26	,875