

## DAFTAR PUSTAKA

- Aboubakar, Njintang, Y. N., Scher, J., & Mbofung, C. M. F. (2008). Physicochemical, thermal properties and microstructure of six varieties of taro (*Colocasia esculenta* L. Schott) flours and starches. *Journal of Food Engineering*, 86(2), 294–305. <https://doi.org/10.1016/j.jfoodeng.2007.10.006>
- Aprianita, A., Purwandari, U., Watson, B., & Vasiljevic, T. (2009). Physico-chemical properties of flours and starches from selected commercial tubers available in Australia. *International Food Research Journal*, 16(4), 507–520.
- Asavasaksakul, S., Tattiyakul, J. and Pradipasena, P. (2005). Chemical and Physical Properties of Taro Starch *Colocasia esculenta* (L.) Schott Grown in Thailand. In *The 3rd Conference on Starch Technology* (pp. 191–197). Bangkok, Thailand: Department of Food Technology, Faculty of Science, Chulalongkorn University, Payathai Road, Patumwan, Bangkok 10330 Thailand.
- Bartholomew, S. C. M. and D. P. (1978). Calcium Nutrition of Taro (*Colocasia esculenta*) (L.) Schoot. In *International Symposium on Tropical Root and Tuber Crops* (pp. 665–670). Honolulu, Hawaii: University of Hawaii.
- Bellettini, I. C., Cristina, A., Luiz, P., Barreto, M., & Soldi, V. (2017). LWT - Food Science and Technology Edible carboxymethyl cellulose films containing natural antioxidant and surfactants: a -tocopherol stability , in vitro release and film properties. *Food Science and Technology*, 77, 21–29. <https://doi.org/10.1016/j.lwt.2016.11.026>
- Bemiller, J. N. (2011). Pasting , paste , and gel properties of starch – hydrocolloid combinations. *Carbohydrate Polymers*, 86(2), 386–423. <https://doi.org/10.1016/j.carbpol.2011.05.064>
- Bird, H. E. M. and M. R. (2014). Gum Arabic Fractionation Using Synthetic Membranes: The Importance Of Fuling. *Food and Bioproducts Processing*, 1–14. <https://doi.org/10.1016/j.fbp.2014.10.008>
- Bong, W. C., Vanhanen, L. P., & Savage, G. P. (2017). Addition of calcium compounds to reduce soluble oxalate in a high oxalate food system. *Food Chemistry*, 221, 54–57. <https://doi.org/10.1016/j.foodchem.2016.10.031>
- Bradbury, J. H., Bradshaw, K., Jealous, W., Holloway, W. D., & Phimpisane, T. (1988). Effect of cooking on nutrient content of tropical root crops from the south pacific. *Journal of the Science of Food and Agriculture*, 43(4), 333–342. <https://doi.org/10.1002/jsfa.2740430406>

- Burey, P., Bhandari, B. R., Howes, T., & Gidley, M. J. (2016). Hydrocolloid Gel Particles: Formation, Characterization, and Application. *Critical Reviews in Food Science and Nutrition*, 48, 361–377. <https://doi.org/10.1080/10408390701347801>
- Cash, M. J., & Caputo, S. J. (2010). *Food Stabilizers, Thickeners and Gelling Agents. Food Stabilizers, Thickeners and Gelling Agents.* Retrieved from <http://blog.ub.ac.id/fawzy/files/2013/07/Food-Stabilizer.pdf#page=66>
- Catherwood, D. J., Savage, G. P., Mason, S. M., Scheffer, J. J. C., & Douglas, J. A. (1989). Oxalate content of cormels of Japanese taro (Colocasia esculenta (L.) Schott) and the effect of cooking. *Journal of Agricultural and Food Chemistry*, 37(2), 147–151. <https://doi.org/10.1016/j.jfca.2005.12.012>
- Chiumarelli, M., & Hubinger, M. D. (2012). Stability, solubility, mechanical and barrier properties of cassava starch e Carnauba wax edible coatings to preserve fresh-cut apples. *Food Hydrocolloids*, 28, 59–67. <https://doi.org/10.1016/j.foodhyd.2011.12.006>
- Chiumarelli, M., & Hubinger, M. D. (2014). Food Hydrocolloids Evaluation of edible films and coatings formulated with cassava starch, glycerol, carnauba wax and stearic acid. *Food Hydrocolloids*, 38, 20–27. <https://doi.org/10.1016/j.foodhyd.2013.11.013>
- Danijela Z. Suput, Vera L. Latic, Senka Z. Popovic, N. M. H. (2015). Edible Films and Coatings – Sources, Properties and Application. *Food and Feed Research*, 42, 11–22. <https://doi.org/10.5937/FFR1501011S>
- Delya Arsi Diova, YS. Darmanto, L. R. (2013). Karakteristik Edible Film Komposit Semirefined Karaginan dari Rumput Laut Eucheuma Cottonii dan Beeswax. *Pengolahan Dan Bioteknologi Hasil Perikanan*, 2(3), 1–10.
- Dewi Artanti Putri, A. C. K. (2013). The Effect of Pretreatment Process for Taro tuber (Colocasia esculenta) Calcium Oxalate Content. In *Engineering International Conference “UNNES Conservation” 2013 Proceeding* (pp. 115–118). Semarang, Indonesia: Department of Chemical Engineering, Engineering Faculty, Diponegoro University, Semarang, Indonesia.
- Durango, A. M., & Andrade, N. J. (2006). Microbiological evaluation of an edible antimicrobial coating on minimally processed carrots. *Food Control*, 17, 336–341. <https://doi.org/10.1016/j.foodcont.2004.10.024>
- Ema susiana, Tesri Maideliza, dan M. (2013). Analysis Morphology of Starch Grains and Crystals in Some Taro. *Jurnal Biologi Universitas Andalas*, 2(4), 249–255.

- Ervika Rahayu N.H., Dini Ariani, Miftakhussolikhah, Maharani P.E., and Y. P. (2017). Chemical Characteristic and Functional Properties of Arenga Starch-Taro ( Colocasia esculanta L .) Flour Noodle with Turmeric Extracts Addition. In *International Symposium on Applied Chemistry (ISAC) 2016* (Vol. 20023). Indonesia: AIP Publishing. <https://doi.org/10.1063/1.4973150>
- Frey-Wyssling, A. (1981). Crystallography of the Two Hydrates of Crystalline Calcium Oxalate in Plants. *American Journal of Botany*, 68, 130–141.
- Garti, N., Madar, Z., Aserin, A., & Sternheim, B. (1997). Fenugreek Galactomannans as Food Emulsifiers. *LWT - Food Sci. Tech.*, 30, 305–311. <https://doi.org/10.1006/fstl.1996.0179>
- Gidley, M. J., & Reid, J. S. G. (2006). Galactomannans and Other Cell Wall Storage Polysaccharides in Seeds. In A. M. Stephen (Ed.), *Food Polysaccharides and Their Applications* (Second, pp. 181–215). London, New York: Taylor and Francis Group, LLC. <https://doi.org/doi:10.1201/9781420015164.ch6>
- Guilbertt, S., Gontards, N., & Cuqs, B. (1995). Technology and Applications of Edible Protective Films. *Packaging Technology and Science*, 8, 339–346.
- Hanani, Z. A. N., Beatty, E., Roos, Y. H., & Morris, M. A. (2013). Development and Characterization of Biodegradable Composite Films Based on Gelatin Derived from Beef, Pork and Fish Sources. *Foods*, 2, 1–17. <https://doi.org/10.3390/foods2010001>
- Harris, P. J., Ferguson, L. R., Roberton, A. M., McKenzie, R. J., & White, J. B. (1991). Cell-wall histochemistry and anatomy of taro (*Colocasia esculenta*). *Australian Journal of Botany*, 39(3), 207–222. <https://doi.org/10.1071/BT9910207>
- Himedialabs. (2017). Product Information Amylum (Starch soluble). Mumbai, India: [Himedialabs.com](http://Himedialabs.com).
- Jane, J., Shen, L., Chen, J., Lim, S., Kasemsuwan, T., & Nip, W. K. (1992). Physical and Chemical Studies of Taro Starches and Flours. *Cereal Chemistry*, 69, 528–535.
- Karimi, A. and B. (2012). Lectins and Their Roles in Pests Control. In P. A. R. Bandani (Ed.), *New Perspectives in Plant Protection* (p. 246). InTech. Retrieved from <http://www.intechopen.com/books/new-perspectives-in-plant-protection/lectins-and-their-roles-in-pest-control>

- Kaur, M., Kaushal, P., & Sandhu, K. S. (2013). Studies on physicochemical and pasting properties of Taro (*Colocasia esculenta* L.) flour in comparison with a cereal, tuber and legume flour. *Journal of Food Science and Technology*, 50(1), 94–100. <https://doi.org/10.1007/s13197-010-0227-6>
- Kaushal, P., Kumar, V., & Sharma, H. K. (2015). Utilization of taro (*Colocasia esculenta*): a review. *Journal of Food Science and Technology*, 52(1), 27–40. <https://doi.org/10.1007/s13197-013-0933-y>
- Koocheki, A., Ghandi, A., Razavi, S. M. A., Mortazavi, S. A., & Vasiljevic, T. (2009). The rheological properties of ketchup as a function of different hydrocolloids and temperature. *International Journal of Food Science and Technology*, 44(3), 596–602. <https://doi.org/10.1111/j.1365-2621.2008.01868.x>
- Kumoro, A. C., Budiyati, C. S., & Retnowati, D. S. (2014). Calcium oxalate reduction during soaking of giant taro (*Alocasia macrorrhiza* (L.) Schott) corm chips in sodium bicarbonate solution. *International Food Research Journal*, 21(4), 1583–1588.
- Kumoro, A. C., Putri, R. D. A., Budiyati, C. S., Retnowati, D. S., & Ratnawati. (2014). Kinetics of Calcium Oxalate Reduction in Taro (*Colocasia Esculenta*) Corm Chips during Treatments Using Baking Soda Solution. *Procedia Chemistry*, 9, 102–112. <https://doi.org/10.1016/j.proche.2014.05.013>
- Lewu, M. N., Adebola, P. O., & Afolayan, A. J. (2010a). Comparative assessment of the nutritional value of commercially available cocoyam and potato tubers in South Africa. *Journal of Food Quality*, 33(4), 461–476. <https://doi.org/10.1111/j.1745-4557.2010.00325.x>
- Lewu, M. N., Adebola, P. O., & Afolayan, A. J. (2010b). Effect of cooking on the mineral contents and anti-nutritional factors in seven accessions of *Colocasia esculenta* (L.) Schott growing in South Africa. *Journal of Food Composition and Analysis*, 23(5), 389–393. <https://doi.org/10.1016/j.jfca.2010.02.006>
- Lin, M. A. P. and S.-J. (2013). The Application of Edible Polymeric Films and Coatings in the Food Industry. *Food Processing and Technology*, 4(2), 1–2. <https://doi.org/10.4172/2157-7110.1000e116>
- Lorenzo, G., Zaritzky, N. E., & Califano, A. N. (1990). Viscoelastic characterization of fluid and gel like food emulsions stabilized with hydrocolloids. *Italian Oral Surgery*, 1(2011), 281–286. <https://doi.org/10.1016/j.profoo.2011.09.044>
- Lorenzo, G., Zaritzky, N. E., & Califano, A. N. (2011). Viscoelastic characterization of fluid and gel like food emulsions stabilized with

- hydrocolloids. *Procedia Food Science*, 1, 281–286. <https://doi.org/10.1016/j.profoo.2011.09.044>
- Maga, J. A. (1992). Taro: composition and food uses. *Food Reviews International*, 8(3), 443–473. <https://doi.org/10.1080/87559129209540948>
- Malone, M. E., Appelqvist, I. A. M., & Norton, I. T. (2003). Oral behaviour of food hydrocolloids and emulsions. Part 2. Taste and aroma release. *Food Hydrocolloids*, 17(6), 775–784. [https://doi.org/10.1016/S0268-005X\(03\)00098-5](https://doi.org/10.1016/S0268-005X(03)00098-5)
- Manchun, S., Nunthanid, J., Limmatvapirat, S., & Sriamornsak, P. (2012). Effect of Ultrasonic Treatment on Physical Properties of Tapioca Starch. *Advanced Materials Research*, 506, 294–297. <https://doi.org/10.4028/www.scientific.net/AMR.506.294>
- Mergedus, A., Kristl, J., Ivancic, A., Sober, A., Sustar, V., Krizan, T., & Lebot, V. (2015). Variation of mineral composition in different parts of taro (*Colocasia esculenta*) corms. *Food Chemistry*, 170, 37–46. <https://doi.org/10.1016/j.foodchem.2014.08.025>
- Milani, J., & Maleki, G. (2012). Hydrocolloids in Food Industry. In *Food Industrial Processes-Methods and Equipment* (p. 418). InTech. <https://doi.org/10.5772/2491>
- Misra, R. S., Sharma, K., & Mishra, A. K. (2008). Phytophthora Leaf Blight of Taro ( *C olocasia esculenta* ) – A Review. *Journal of Plant Science and Biotechnology*, 2, 55–63.
- Mittal, S. A. and P. (2014). Recent Trends in Edible Films and Coatings for Fresh and Minimally Processed Fruits. *The International Journal of Science & Technolodge*, 2(12), 239–244.
- Nand, A. V, Charan, R. P., Rohindra, D., & Khurma, J. R. (2008). Isolation and properties of starch from some local cultivars of cassava and taro in Fiji. *The South Pacific Journal of Natural Science*, 26, 45–48.
- O. Skurys. (2010). *Food Hydrocolloid Edible Films and Coatings*. (O. Skurys, Ed.). Santiago, Chile. Phone: Nova Science Publishers.
- Oecd SIDS. (2002). Sodium Bicarbonate OECD SIDS. Boston, USA: UNEP PUBLICATIONS.
- Onwueme, I. (1999). *Taro Cultivation in Asia and In The Pacific*. Food and Agriculture Organization of The United Nations Regional Office for Asia and The Pacific. Bangkok: RAP PUBLICATIONS.

- Otang, W. M., Grierson, D. S., & Afolayan, A. J. (2014). A survey of plants responsible for causing irritant contact dermatitis in the Amathole district, eastern cape, South Africa. *Journal of Ethnopharmacology*, 157, 274–284. <https://doi.org/10.1016/j.jep.2014.10.002>
- Patrícia Ribeiro Pereira, Eduardo Mere Del Aguila, Maurício Afonso Verícimo, R. B. Z., Paschoalin, V. M. F., & Silva, J. T. (2014). Purification and characterization of the lectin from taro (*coccosia esculenta*) and its effect on mouse splenocyte proliferation in vitro and in vivo. *Protein Journal*, 33(1), 92–99. <https://doi.org/10.1007/s10930-013-9541-y>
- Pereira, P. R., Meagher, J. L., Winter, H. C., Goldstein, I. J., Paschoalin, V. M., Silva, J. T., & Stuckey, J. A. (2016). High-resolution crystal structures of *Coccosia esculenta* tarin lectin. *Glycobiology*, 1–7. <https://doi.org/10.1093/glycob/cww083>
- Pramodrao, K. S., & Riar, C. S. (2013). Food Hydrocolloids Comparative study of effect of modifi cation with ionic gums and dry heating on the physicochemical characteristic of potato , sweet potato and taro starches. *Food Hydrocolloids*. <https://doi.org/10.1016/j.foodhyd.2013.08.006>
- Prana, M. S. R. I. (2007). Study on Flowering Biology of Taro (*Coccosia esculenta* (L.) Schott.). *Biodiversitas*, 8(1), 63–66.
- Rosell, C. M., Rojas, J. A., & Benedito de Barber, C. (2001). Influence of hydrocolloids on dough rheology and bread quality. *Food Hydrocolloids*, 15(1), 75–81. [https://doi.org/10.1016/S0268-005X\(00\)00054-0](https://doi.org/10.1016/S0268-005X(00)00054-0)
- Saha, D., & Bhattacharya, S. (2010). Hydrocolloids as thickening and gelling agents in food: a critical review. *Journal of Food Science and Technology*, 47(6), 587–597. <https://doi.org/10.1007/s13197-010-0162-6>
- Santoso, B., Pitayati, P. A., & Pambayun, R. (2013). The use of Carrageenan and Gum Arabic for Hidrocolloid Based Edible Film. *Agritech*, 33(2), 140–145.
- Savage, N. and. (1999). Oxalate Content of Foods and its Effect on Humans. *Asia Pacific Journal Clinic and Nutrition*, 8(1), 64–74.
- Sefa-Dedeh, S., & Agyir-Sackey, E. K. (2004). Chemical composition and the effect of processing on oxalate content of cocoyam *Xanthosoma sagittifolium* and *Coccosia esculenta* cormels. *Food Chemistry*, 85(4), 479–487. [https://doi.org/10.1016/S0308-8146\(02\)00244-3](https://doi.org/10.1016/S0308-8146(02)00244-3)
- Setiani, W., Sudiarti, T., & Rahmidar, L. (2013). Preparasi Dan Karakterisasi Edible Film Dari Poliblend Pati Sukun-Kitosan. *Jurnal Kimia Valensi*, 3(2), 100–109.

Shah, U., Naqash, F., Gani, A., & Masoodi, F. A. (2016). Art and Science behind Modified Starch Edible Films and Coatings: A Review. *Comprehensive Reviews in Food Science and Food Safety*, 15, 568–580. <https://doi.org/10.1111/1541-4337.12197>

Shaikh, S. F., Mane, R. S., Min, B. K., & Hwang, Y. J. (2016). D-Sorbitol-Induced Phase Control of TiO<sub>2</sub> Nanoparticles and its Application for Dye-Sensitized Solar Cells. *Nature Publishing Group*, 6, 1–10. <https://doi.org/10.1038/srep20103>

Sinaga, R. F., Ginting, G. M., Hasibuan, R., Kimia, D. T., Teknik, F., & Utara, U. S. (2014). Pengaruh Penambahan Gliserol terhadap Sifat Kekuatan Tarik dan Pemanjangan Saat Putus Bioplastik dari Pati Umbi Talas. *Jurnal Teknik Kimia Universitas Sumatera Utara*, 3(2), 19–24.

Slaughter, S. L., Ellis, P. R., Jackson, E. C., & Butterworth, P. J. (2002). The effect of guar galactomannan and water availability during hydrothermal processing on the hydrolysis of starch catalysed by pancreatic α-amylase. *Biachimica et Biophysica Acta*, 1571, 55–63.

Sri Wahyu Murni, Harso Pawignyo, Desi Widyawati, N. S. (2013). Pembuatan Edible Film dari Tepung Jagung (*Zea Mays L.*) dan Kitosan. In *Seminar Nasional Teknik Kimia “Kejuangan” Pengembangan Teknologi Kimia untuk Pengolahan Sumber Daya Alam Indonesia*. Yogyakarta, Indonesia: Program Studi Teknik Kimia, FTI, UPN “Veteran” Yogyakarta. <https://doi.org/10.6066/jtip.2013.24.1.48>

Srikaeo, K., Mingyai, S., & Sopade, P. A. (2011). Physicochemical properties, resistant starch content and enzymatic digestibility of unripe banana, edible canna, taro flours and their rice noodle products. *International Journal of Food Science and Technology*, 46(10), 2111–2117. <https://doi.org/10.1111/j.1365-2621.2011.02724.x>

Strauss, M. S., Stephens, G. C., Gonzales, C. J., & Arditti, J. (1980). Genetic Variability in Taro , *Colocasia esculenta* ( L .) Schott ( Araceae ). *Annals of Botany Company*, 45, 429–437.

Temesgen, M. (2015). Nutritional Potential , Health and Food Security Benefits of Taro *Colocasia Esculenta* ( L .): A Review. *Food Science and Quality Management*, 36, 23–31.

Vajravijayan, S., Pletnev, S., Pletnev, V. Z., Nandhagopal, N., & Gunasekaran, K. (2016). Structural analysis of β-prism lectin from *Colocasia esculenta* (L.) S chott. *International Journal of Biological Macromolecules*, 91, 518–523. <https://doi.org/10.1016/j.ijbiomac.2016.05.048>

Van Damme, E., Coossens, K., Smeets, K., Van Leuven, F., Verhaert, P., & J. Peumans, W. (1995). The Major Tuber Storage Protein of Araceae Species 1s a Lectin, (107), 1147–1158.

Vilgis, T. A. (2012). Hydrocolloids between soft matter and taste: Culinary polymer physics. *International Journal of Gastronomy and Food Science*, 1(1), 46–53. <https://doi.org/10.1016/j.ijgfs.2011.11.012>

Wang, J.-K. S. H. (1983). *Taro A Review Of Colocasia Esculenta and Its Potensial*. (J.-K. S. H. Wang, Ed.). Hawai, USA: University of Hawaii Press.

Warkoyo , Budi Rahardjo , Djagal Wiseso Marseno, J. N. W. K. (2014). Physical, Mechanical and Barrier Properties of Xanthosoma sagittifolium Starch-Based Edible Film Incorporated with Potassium Sorbate. *Agritech*, 34(1), 72–81.

Wüstenberg, T. (2015). General Overview of Food Hydrocolloids. *Cellulose and Cellulose Derivatives in the Food Industry Fundamentals and Applications*, 1–68. Retrieved from [http://media.johnwiley.com.au/product\\_data/excerpt/8X/35273375/35273375\\_8X-218.pdf](http://media.johnwiley.com.au/product_data/excerpt/8X/35273375/35273375_8X-218.pdf)

Young-Joo Seo, Satsuki Une, Ikuyo Tsukamoto, and M. M. (1989). The Effect of Lectin from Taro Tuber (*Colocasia antiquorum*) Given by Force-Feeding on the Growth of Mice. *Journal Nutrition Science Vitaminol*, 36, 277–285.